


Oxytarm™ and fibromyalgia

Results after 8 weeks. 17 person started out, 12 was controlled after 8 weeks and 10 after 16 weeks.

1. Question: How much general body pain have you had?

On a visual analogue scale from 0 to 10 where 0 is strong pain and 10 is no pain.


Before the study:	score 3,5
After 8 weeks:	score 5,5
After 16 weeks:	score 6,5



A 20% reduction in pains after 2 months
A 30% reduction in pains after 4 months

2. Question: How much difficulty have you had with your daily activities?
0 is "could not do any activity" and 10 is no difficulty.

Before the study:	score 5,5
After 8 weeks:	score 7,5
After 16 weeks:	score 8,0




20% improvement of daily activity after 2 months
25% improvement of daily activity after 4 months


3. Question: What is the hardest physical activity you could do for at least 2 minutes?

0 is very light (slow walking), 10 is very strong (fast running).

Before the study:	score 4
After 8 weeks:	score 5,5
After 16 weeks:	score 6

15% increase in physical activity after 2 months
20% increase in physical activity after 4 months





4. Question: How would you classify your general health condition?
0 is poor and 10 is excellent

Before the study: score 3
After 8 weeks: score 6
After 16 weeks: score 6,5

30% improvement in general health condition after 2 months
35% improvement in general health condition after 4 months

